

Mahlet Abayneh
Kullehe Haddis Yeshanew
Samuel Lulseged Yidengitu
Redeat Workneh Tadesse
Afolayan Ayomide
Erika M. Edwards
Danielle E. Y. Ehret
Pamela Henderson
Loveness Nya Kwima
Benenia Muzuva
John Baptist Nkuranga
Helina Selam
Elizabeth Siraha
Alexander G. Stevenson
Misrak Tadesse
Gregory C. Valentine

<https://dx.doi.org/10.4314/jan.v3i3.5>

Received: 1st July 2025

Accepted: 7th July 2025

Mahlet Abayneh (✉)
 Redeat Workneh Tadesse
 Samuel Lulseged Yidengitu
 St Paul's Hospital Millennium
 Medical College, Addis Ababa,
 Ethiopia
 Email:
 mahlet.abayneh@sphmmc.edu.et

Kullehe Haddis Yeshanew
 Tikur Anbessa Specialized Hospital,
 Addis Ababa, Ethiopia

Afolayan Ayomide
 Federal Teaching Hospital,
 Ido-Ekiti, Nigeria

Erika M. Edwards
 Danielle E. Y. Ehret
 Vermont Oxford Network and
 University of Vermont,
 Burlington, Vermont, USA

Pamela Henderson
 Alexander G. Stevenson
 African Neonatal Network, Kigali,
 Rwanda and Harare, Zimbabwe



Nutrition practices of neonatal intensive care units in the African Neonatal Network

Abstract: *Background:* Optimal and adequate nutrition provision and standard nutritional practice is mandatory for critically ill infants in the neonatal intensive care unit. Poor nutrition in early life is associated with problems that increase risk of mortality and morbidity in preterm newborns. Preterm infants are at high risk of having inadequate nutrition in low-resource settings due to lack of resources and trained health workers.

Methods: Fourteen hospitals in the African Neonatal Network responded to an annual facility survey and a health facility survey co-developed by faculty in the African Neonatal Network and Vermont Oxford Network. All analyses use descriptive statistics.

Results: Almost all the units (94%) had guidelines for initiation and advancement of feeding. Ten hospitals (71%) reported complying with Baby-Friendly Hospital initiatives. Among the 14 hospitals, only one (7%) had access to pasteurized donor milk. There was significant variation in use of breast milk fortifiers; only three (21%) hospitals reported using it consistently for more than 90% of their preterm newborns. There was also variation in human resources related to neonatal NICU nutrition with only four (29%) of the hospi-

tals having a dedicated dietitian and one with 24/7 access to a lactation consultant.

Conclusion: Dedicated dietitian or lactation support team were limited in the network hospitals. There was only one hospital with access to pasteurized donor milk, and use of breast milk fortifiers was not optimal. Optimising nutrition should be prioritised to improve opportunities for infants to go home meeting growth standards.

Keywords: Infant, Premature; Infant, Newborn; Nutritional Support; Neonatal Intensive Care Units; Lactation Support; Diet, Infant; Milk, Human; Human Milk Banking; Africa South of the Sahara; Global Health

Résumé: *Contexte:* Une nutrition optimale et adéquate, ainsi que des pratiques nutritionnelles standardisées, sont indispensables pour les nourrissons gravement malades en unité de soins intensifs néonataux (USIN). Une mauvaise nutrition au début de la vie est associée à des complications qui augmentent le risque de mortalité et de morbidité chez les nouveau-nés prématurés. Dans les contextes à faibles ressources, ces nourrissons sont particulièrement exposés à

Loveness Nya Kwima,
Benenia Muzuva
Mbuya Nehanda Maternity Hospital,
Harare, Zimbabwe

John Baptist Nkuranga
University of Rwanda/African
Health Sciences University, Kigali,
Rwanda

Helina Selam
Vermont Oxford Network,
Burlington, Vermont, USA

Elizabeth Siraha
Great Zimbabwe University,
Masvingo, Zimbabwe

Misrak Tadesse
Vermont Oxford Network and Johns
Hopkins University School of
Medicine, Baltimore, Maryland,
USA

Gregory C. Valentine
University of Washington,
Seattle, Washington, USA

une nutrition inadéquate en raison
du manque de ressources et de
personnel de santé formé.

Méthodes: Quatorze hôpitaux du
Réseau Néonatal Africain ont ré-
pondu à une enquête annuelle sur
les établissements et à une enquête
sur les structures de santé, co-
développées par des membres du
Réseau Néonatal Africain et du
Vermont Oxford Network. Toutes
les analyses ont été réalisées à
l'aide de statistiques descriptives.

Résultats: Presque toutes les uni-
tés (94%) disposaient de proto-
coles pour l'initiation et la pro-
gression de l'alimentation. Dix
hôpitaux (71%) déclaraient se
conformer à l'initiative des Hôpi-
taux Amis des Bébé. Parmi les 14
hôpitaux, un seul (7%) avait accès
à du lait de donneuse pasteurisé.
Une variation importante a été
observée dans l'utilisation des
fortifiants du lait maternel: seule-
ment trois hôpitaux (21%) dé-

claraient les utiliser systématique-
ment pour plus de 90% de leurs
nouveau-nés prématurés. La dis-
ponibilité en ressources humaines
spécialisées dans la nutrition neo-
natal était également variable :
seuls quatre hôpitaux (29%) dis-
posaient d'un·e diététicien·ne
dédié·e, et un seul avait un accès
24 h/24 à un·e consultant·e en lac-
tation.

Conclusion: Le soutien nutrition-
nel spécialisé par un·e diététicien·
ne ou une équipe de lactation était
limité dans les hôpitaux du réseau.
Un seul hôpital avait accès à du
lait de donneuse pasteurisé, et
l'utilisation des fortifiants du lait
maternel restait sous-optimale.
L'optimisation de la nutrition doit
être une priorité afin de maximiser
les chances pour les nourrissons
de sortir de l'hôpital en atteignant
les normes de croissance.

Introduction

Critically ill infants in neonatal intensive care units re-
quire appropriate nutrition for survival, growth and de-
velopment.¹ Inadequate nutrition in the first postnatal
week results in poor growth, especially in very low birth
weight (VLBW) infants.² Faltering growth of VLBW
infants during postnatal hospitalization is associated
with impaired neurodevelopmental outcomes, increased
risk of retinopathy of prematurity, persistent growth
deficits and increased risk for cardiovascular disease in
adulthood.^{2,3}

Early and adequate nutrition for critically ill infants im-
proves growth, shortens length of hospital stay, and de-
creases development of late onset neonatal sepsis, necro-
tising enterocolitis, and chronic lung disease, especially
in preterm infants.⁴ Current recommendations suggest
that the goal of postnatal nutrition in preterm newborns
should be to replicate the needs of the foetus in the same
gestational age.⁵ Poor or inadequate nutrition postnatally
resulting in extrauterine growth restriction (EUGR) has
commonly been considered as growth values <10th per-
centile of expected intrauterine growth or as a decrease
in z-scores between birth and a defined postmenstrual
age (PMA).⁶

Optimizing nutrition for these vulnerable small and/or
sick newborns in low-resourced settings worldwide is
challenging given the lack of human and physical re-
sources including availability of parenteral nutrition and
its macronutrients. Thus, it is not surprising that publica-
tions have demonstrated high rates of EUGR among
preterm newborns.^{7,8} For example, among preterm new-

borns, 86.2% discharged from Ethiopian NICUs and up
to 85.7% in South Africa had EUGR.^{7,8}

Additionally, ensuring ideal growth in preterm new-
borns requires optimization of both parenteral fluids (for
those that require intravenous fluids) and enteral feeds.
Timely initiation of enteral feeds, ideally via breastfeed-
ing within 1 hour after birth, is a current recommenda-
tion by the World Health Organization.⁹ Furthermore,
early initiation and advancement of enteral feeding is
recommended for VLBW infants.¹⁰ However, feeding
initiation and advancement is a challenge in VLBW in-
fants due to acute illnesses and functional immaturity of
gastrointestinal system, along with limited availability
of total parenteral nutrition, breastfeeding support, and
donor milk.¹¹ Having standard feeding protocols, using
mother's own milk, and engagement of a multidisciplinary
team are recommended methods to reduce feeding
challenges in the neonatal intensive care unit.¹²

There is paucity of data in low- and middle-income
countries on nutritional practices and nutritional out-
comes of preterm newborns discharged from NICUs in
these settings. There has never been a review of the cur-
rent nutritional practices within NICUs within the Afri-
can continent. The African Neonatal Network (ANN)
includes 14 hospitals across five countries: Ethiopia,
Nigeria, Rwanda, Uganda, and Zimbabwe. Leveraging
the ANN, we aimed to assess the nutritional practices
and discharge weight of preterm newborns across sub-
Saharan Africa to provide an overview of current prac-
tices and identify areas of potential improvements to
optimize nutritional outcomes for small and sick new-
borns in these settings.

Methods

Data collection was conducted in 14 African Neonatal Network (ANN) member hospitals across five countries: Ethiopia, Nigeria, Rwanda, Uganda, and Zimbabwe.

Vermont Oxford Network (VON) conducts an annual survey for members that was co-developed with ANN faculty members, which includes information on the hospital setting, number of beds and admissions, staffing, obstetric service, follow-up clinic, resuscitation and essential newborn care, transfers and transport, family-centred care, services provided by the neonatal unit, guidelines in the neonatal unit, quality assurance/continuous quality improvement, and level of neonatal care. Participation in the membership survey is mandatory. The responses used for this manuscript are from 2023.

In October 2023, the ANN conducted a health facility assessment to collect more detailed information on buildings and facilities, medications, diagnostics and consumables, equipment, staffing, governance, thermal regulation and foetal transition, nutrition, family-centred care and kangaroo mother care, infection prevention and control, and perceived priorities.

Tables of hospital-level measures include data from both the membership survey and the health facility assessment. All analyses are descriptive. The collaborative QI project and subsequent assessments received individual and hospital institutional research and ethics review approvals at the start of the collaborative and learning initiative.

Results

Among the 14 hospitals, four (29%) had dedicated paediatric nutritionists or dietitians. Ten hospitals (71%) had nurses trained on lactation support. However, only one hospital reported that a lactation consultant was available 24 hours per day, two (14%) hospitals reported one was available most days during work hours, three (21%) reported one was occasionally available but not on the ward every day, and eight (38%) reported that lactation support was not generally available.

Ten (71%) hospitals complied with the Baby Friendly Hospital initiative, and guidelines for initiation of breastfeeding and advancement of feeding were used consistently in 13 (93%) hospitals. All hospitals reported that weight scales and nasogastric or orogastric tubes were consistently available. Ten (71%) reported the availability of refrigerators for expressed milk, sterilized cups, and electrolyte diagnostics but only four (29%) had total parenteral nutrition (Table 1).

Table 1: Supplies consistently available at 14 African Neonatal Network member hospitals

	#	%
Weight scale	14	100
Nasogastric or orogastric tubes for feeding	14	100
Hypoglycemia diagnostics	11	79
Refrigerator for expressed milk	10	71
Sterilizable cups for complementary feeding	10	71
Nursing staff trained in supporting lactation	10	71
Electrolyte diagnostics	10	71
Infusion pumps	9	64
Term and preterm formula on formulary	7	50
Human milk fortifier	4	29
Total parenteral nutrition	4	29

A dedicated area for preparing or mixing milk/feeds was found in eight (57%) and seven (50%) had a separate space within the neonatal unit to allow mothers to express milk privately (Table 2). Only one hospital provided electrical breast milk pumps, and none provided mothers with manual breast pumps. Ten hospitals (71%) provided containers for mothers to express milk. Only one hospital had regular access to donated and pasteurized breast milk from a formal breast milk bank. However, two units utilized donated breast milk from persons other than mother but not through a formal breast milk bank.

Breastfeeding counselling for mothers during the antenatal care was practiced more than 60% of the time at 10 (71%) hospitals, while three (21%) hospitals had no practice of antenatal counselling on breastfeeding and two hospitals (14%) did not know whether antenatal counselling was available. Five hospitals (36%) reported that over 90% of mothers received lactation encouragement, support and education in the immediate postnatal period, while seven hospitals (50%) reported that 41-90% of mothers received such help and two hospitals reported that 40% or fewer mothers received such help. Most hospitals provided regular vitamin D fortification (64%) and iron supplementation (71%) to infants born <32 weeks, but only one hospital (7%) had zinc supplementation (Table 3).

Eight hospitals (57%) reported providing special preterm formula for preterm infants when clinically indicated. Three (21%) reported that some infants receive preterm formula while some receive term formula. One (7%) hospital reported that most preterm infants received term formula. Provision of preterm formula was reported as unknown by two (14%) hospitals.

Two hospitals reported that >90% of nurses and physicians received training on breastfeeding support after qualification while five (36%) hospitals reported that no more than 10% of nurses received training, and six (43%) reported that no more than 10% of physicians received training (Table 4).

Table 2: Institutional support for breastfeeding at 14 African Neonatal Network member hospitals

	#	%
Refrigerated storage space for mothers to store expressed milk	10	71
Hospital provides containers for mothers to express milk	10	71
Dedicated area (not at bedside) for preparing or mixing milk/feeds	8	57
Separate space within neonatal unit that allows mothers to express milk privately	7	50
Unit utilizes donated breast milk from person other than mother but not from formal breast milk bank	2	14
Regular access to donated and pasteurized breast milk from formal breast milk bank	1	7
Hospital provides electric breast pumps	1	7
Hospital provides manual breast pumps	0	0

Table 3: Hospital fortification and supplementation practices for infants at 14 African Neonatal Network member hospitals

	Preterm infants receiving breast milk fortification		<32 weeks receiving regular vitamin D fortification		<30 weeks receiving iron supplementation		<32 weeks receiving zinc supplementation		<32 weeks receiving suitable multivitamin	
	#	%	#	%	#	%	#	%	#	%
None	2	14	1	7			9	64	5	35
≤10%	3	21	1	7	1	7	1	7		
11% to 40%	2	14			1	7				
41% to 60%	2	14							1	7
61% to 90%	2	14	3	21	2	14	1	7	1	7
>90%	3	21	9	64	10	71	1	7	7	50
Unknown							2	14		

Table 4: Percentage of staff trained on breastfeeding support after qualification at 14 African Neonatal Network member hospitals

	Nurses		Physicians	
	#	%	#	%
≤ 10%	5	36	6	43
11%-40%	3	21	2	14
41%-60%	1	7	3	21
61%-90%	3	21	1	7
>90	2	14	2	14

Discussion

This survey evaluated the current nutritional practices of 14 neonatal intensive care units in five countries that are members of the African Neonatal Network. Multidisciplinary team involvement, including dietitians in American NICUs showed improvement in weight gain.¹³ However only 4 (29%) NICUs had dedicated dietitians in the network hospitals.

Ten hospitals reported having nurses trained in lactation support. Training of lactation support for NICU nurses has demonstrated improvement in lactation knowledge and attitudes, creating a lactation supportive atmosphere in the NICU and improving breastfeeding rates.^{14,15} Although counselling of mothers during pregnancy about breastfeeding has improved early initiation and

exclusive breastfeeding rate,^{16,17} only four hospitals among the 14 had mothers counseled on exclusive breastfeeding during antenatal care.

Baby-Friendly Hospital initiatives practiced by 71% of network hospitals empower early and continued breastfeeding, skin-to-skin, and kangaroo mother care practices. Nutritional guidelines help provide clinicians with standard of feeding practices. Our findings demonstrated that 93% of the participating NICUs are using guidelines for feeding. Standardized feeding and nutrition guidelines and protocols have demonstrated multiple benefits including lesser use of parenteral nutrition, facilitate early enteral nutrition, improved weight gain and reduced rates of necrotising enterocolitis.^{18,19}

Only one hospital reported practicing pasteurized donor milk for admitted infants, St Francis Hospital Nsambya.^[20] All other units reported using formula milk if breast milk is not available from the mother. Early initiation of feeding with mother's breast milk is the recommended feeding for all newborn, including VLBW babies. In scenarios where breast milk is not available, pasteurized donor breast milk is a superior alternative to infant formula or any other breast milk substitute.¹⁰ At three hospitals, more than 90% of qualifying infants received breast milk fortifier; however, three hospitals have never used it. Breast milk fortification for infants on mothers

or donor breast milk is recommended for all very low birth weight infants.^{10,21}

Conclusions

There is significant variation among network hospitals on nutritional practices and availability of staff to support nutrition in the NICU. Many potential areas for better practices were identified, including improving the number of specialist dietitians, improving lactation and nutrition training for doctors and nurses and improving access to both fortification and donated breast milk.

Acknowledgments

We are indebted to our colleagues at the following hospitals who submit data to VON on behalf of infants and their families: St. Paul's Millennium Medical College,

Addis Ababa, Ethiopia; Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia; Tirunesh Beijing Hospital, Addis Ababa, Ethiopia; Assosa Hospital, Asosa, Ethiopia; Hawassa Referral Hospital, Awassa, Ethiopia; Ayder Hospital, Mekelle, Ethiopia; Sacred Heart Hospital, Abeokuta, Nigeria; Federal Teaching Hospital Ido-Ekiti, Ekiti, Nigeria; King Faisal Hospital, Kigali, Rwanda; Mengo Teaching Hospital, Kampala, Uganda; St Francis Nsambya Hospital, Kampala, Uganda; Lubaga Hospital, Kampala, Uganda; Murambinda Mission Hospital, Harare, Zimbabwe; Neocare Baby Hospital, Harare, Zimbabwe.

Funding: Bill and Melinda Gates Foundation INV-042791

References

- Pollack MM, Ruttimann UE, Wiley JS. Nutritional Depletions in Critically Ill Children: Associations with Physiologic Instability and Increased Quantity of Care. *J Parenter Enteral Nutr.* 1985;9(3):309–13.
- Maas C, Poets CF, Franz AR. Avoiding postnatal undernutrition of VLBW infants during neonatal intensive care: evidence and personal view in the absence of evidence. *Arch Dis Child Fetal Neonatal Ed.* 2015;100(1):F76–81.
- Meier PP, Patel AL, Bigger HR, Chen Y, Johnson TJ, Rossman B, et al. Human milk feedings in the neonatal intensive care unit. In: Rajendram R, Preedy VR, Patel VB, editors. *Diet and Nutrition in Critical Care*. Springer New York; 2015. p. 807–22.
- McCallie KR, Lee HC, Mayer O, Cohen RS, Hintz SR, Rhine WD. Improved outcomes with a standardized feeding protocol for very low birth weight infants. *J Perinatol.* 2011;31(S1):S61–7.
- Kleinman RE, Greer FR. *Pediatric Nutrition*. 8th ed. American Academy of Pediatrics Itasca, IL; 2020.
- Ofek Shlomai N, Reichman B, Lerner-Geva L, Boyko V, Bar-Oz B. Population-based study shows improved postnatal growth in preterm very-low-birthweight infants between 1995 and 2010. *Acta Paediatr.* 2014;103(5):498–503.
- Gidi NW, Goldenberg RL, Nigussie AK, et al. Incidence and associated factors of extrauterine growth restriction (EUGR) in preterm infants, a cross-sectional study in selected NICUs in Ethiopia. *BMJ Paediatr Open* 2020;4:e000765.
- Mosidi, LND, van Niekerk A, Coetzee M. 2024. Change in the prevalence of extra-uterine growth restriction in very low birthweight infants, following the introduction of a written nutrition protocol, in a tertiary neonatal unit. *South Afr J Clin Nutr.* 2024;37(3):140–47.
- WHO/UNICEF. Global strategy on infant and young child feeding. Geneva: World Health Organization; 2003.
- WHO. Recommendations for care of the preterm or low birth weight infant. 1st ed. Geneva: World Health Organization; 2022.
- Gephart SM, Hanson C, Wetzel CM, Fleiner M, Umberger E, Martin L, et al. NEC-zero recommendations from scoping review of evidence to prevent and foster timely recognition of necrotizing enterocolitis. *Matern Health Neonatol Perinatol.* 2017 Dec 18;3:23.
- American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.) Board of Directors. Clinical guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients, 2009. *J Parenter Enteral Nutr.* 2009;33(3):255.
- Sneve J, Kattelman K, Ren C, Stevens DC. Implementation of a multidisciplinary team that includes a registered dietitian in a neonatal intensive care unit improved nutrition outcomes. *Nutr Clin Pract.* 2008-2009;23(6):630-4.
- Bernaix LW, Schmidt CA, Arrizola M, Iovinelli D, Medina-Poelinez C. Success of a lactation education program on NICU nurses' knowledge and attitudes. *J Obstet Gynecol Neonatal Nurs.* 2008;37(4):436-45.

15. Maastrup R, Rom AL, Walloe S, Sandfeld HB, Kronborg H. Improved exclusive breastfeeding rates in preterm infants after a neonatal nurse training program focusing on six breastfeeding-supportive clinical practices. *PLoS One*. 2021;16(2):e0245273.
16. Olsen SL, Park ND, Tracy K, Younger D, Anderson B. Implementing standardized feeding guidelines, challenges, and results. *Neonatal Netw*. 2018;37(4):218-223.
17. Wittwer A, Hascoët JM. Impact of introducing a standardized nutrition protocol on very premature infants' growth and morbidity. *PLoS One*. 2020 May 21;15(5):e0232659.
18. Sehhatie FS, Mirghafourvand M, Havizari S. Effect of prenatal counseling on exclusive breastfeeding frequency and infant weight gain in mothers with previous unsuccessful breastfeeding: a randomized controlled clinical trial. *J Matern Fetal Neonatal Med*. 2020 Nov;33(21):3571-3578.
19. Koli H, Nair M, Shah M. (2021). Impact of structured pre-natal counselling on early initiation of breastfeeding and adherence to exclusive breastfeeding: a comparative interventional study. *Int J Reprod Contracept Obstet Gynecol*. 2021;10:2699.
20. Nakibuuka V, Kainza J, Nansiima R, Nalunga S, Nazziwa R, Mponye H, et al. Setting up the first human milk bank in Uganda: a success story from Nsambya hospital. *Front Nutr*. 2024;10:1275877.
21. Moreira DH, Gregory SB, Younge NE. Human milk fortification and use of infant formulas to support growth in the neonatal intensive care unit. *Nutr in Clin Prac*. 2023;38(S2).
22. Stevenson AG, Abayneh M, Bolaji O, et al. Characteristics, interventions, and status of infants discharged from 14 hospitals in the African Neonatal Network, 2024. *J Afric Neonatal*. 2025;3(3):68-77
23. Areprekumor T ebi, Ezech BU, Madjemu RP, Okocha EL. Prevalence and determinants of exclusive breastfeeding among infants after discharge from a neonatal unit in South-South Nigeria. *Int J Trop Dis Health*. 2024;45(12):85-101.
24. Belfort MB, Edwards EM, Greenberg LT, Parker MG, Horbar JD. Diet, weight gain, and head growth in hospitalized U.S. very preterm infants: a 10-year observational study. *Am J Clin Nutr*. 2019;109(5):1373-1379.
25. Fang L, Wu L, Han S, Chen X, Yu Z. Quality improvement to increase breastfeeding in preterm infants: Systematic review and meta-analysis. *Front Pediatr*. 2021 Jun 10;9:681341.